



Reichenbach, L. F., Sobri, A. A., Zaccai, N. R., Agnew, C., Burton, N., Eperon, L. P., de Ornellas, S., Eperon, I. C., Brady, R. L., & Burley, G. A. (2016). Structural Basis of the Mispairing of an Artificially Expanded Genetic Information System. *Chem*, 1(6), 946-958.  
<https://doi.org/10.1016/j.chempr.2016.11.009>

Peer reviewed version

Link to published version (if available):  
[10.1016/j.chempr.2016.11.009](https://doi.org/10.1016/j.chempr.2016.11.009)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cell Press at <http://www.sciencedirect.com/science/article/pii/S2451929416302364> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>




**A** Watson-Crick G-C base-pair**B** AEGIS P-Z base-pair (Benner)**C This work:** Structural basis of pH-dependent mis-pairing of Z with G

Figure 1. Base-pairs of (A) naturally-occurring G-C, (B) P-Z and (C) the pH-dependent G-Z mispair.




Figure 2. (A) Overlay of ODN1 (magenta), ODN2 (orange), ODN3a (green) and ODN3b (cyan) crystal structures. Base-pairs for the 4-position (B) G-C (ODN1), (C) P-Z (ODN2 at pH 7.0), (D) G-Z (ODN3a at pH 7.8), and (E) G-Z (ODN3b at pH 8.5). Hydrogen-bond distances shown in Ångströms. The  $2F_o - F_c$  electron density is contoured at  $1.0\sigma$ .




Figure 3. Comparative analysis of duplex stability and conformation of ODN1-4. (a) UV-vis melting temperatures of ODN1-3 duplexes between pH 5.5 and 9.5 measured at 260 nm. (See also Fig. S9-28, Table S2-3) (b) CD spectra of ODN1-3 at pH 7.5 exhibit characteristic right-handed B-type DNA spectra. (See also Fig. S29-34)




Figure 4. The effect of pH on PCR amplification of a DNA template containing a single P-Z base pair compared to a DNA template containing no P and Z nucleotides. Each PCR was conducted in cacodylate buffer. (See also Fig. 35-37)





Figure 5. Putative resonance structures of the Z anion.

Table 1. Schematic representation of a purine-pyrimidine base-pair in position 4.



	C-1'-C-1' Distance (Å)	N-9-C-1'-C-1' Angle (°)	N-1(C-1)-C-1'-C-1' Angle (°)	Angle of C-1'-C-1' vector (relative to ODN1) (°)
ODN1 <sup>a</sup>	10.4	56.5	55.9	0
ODN2	10.8	55.5	52.3	2.3
ODN3a	10.7	44.5	63.0	9.6
ODN3b	10.8	57.5	54.7	1.0
ODN4 <sup>b</sup>	10.4	42.5	69.9	14.0

Table legend.

<sup>a</sup>PDB ID 463D<sup>b</sup>PDB ID 113D